

<b>PHP_2.01.35      Paraspinal Surface Electromyography to Evaluate and Monitor Back Pain</b>			
<b>Original Policy Date:</b>	December 1, 2025	<b>Effective Date:</b>	December 1, 2025
<b>Section:</b>	2.0 Medicine	<b>Page:</b>	Page 1 of 10

## State Guidelines

As of the publication of this policy, there are no applicable Medi-Cal guidelines (Provider Manual or All Plan Letter). Please refer to the Policy Statement section below.

## Policy Statement

In the absence of any State Guidelines, please refer to the criteria below.

- I. Paraspinal surface electromyography (SEMG) is considered **investigational** as a technique to diagnose or monitor back pain.

## Policy Guidelines

### Coding

See the [Codes table](#) for details.

## Description

A noninvasive procedure that records the summation of muscle electrical activity, paraspinal surface electromyography (SEMG) has been investigated as a technique to evaluate the physiologic functioning of the back. Additionally, this procedure has been studied as a technique to evaluate abnormal patterns of electrical activity in the paraspinal muscles in individuals with back pain symptoms, such as spasm, tenderness, limited range of motion, or postural disorders.

### Summary of Evidence

For individuals who have back pain who receive paraspinal surface electromyography (SEMG) for evaluation and monitoring, the evidence includes several nonrandomized studies on using findings to classify back pain. Relevant outcomes are test accuracy and validity, symptoms, functional outcomes, quality of life, and resource utilization. There have been no studies directly comparing SEMG with other noninvasive techniques for evaluating back pain, and standard criteria for normal and abnormal SEMG measurements have not been determined. Surface electromyography has been proposed as a noninvasive technique providing objective measurements that would inform treatment decisions in patients with back pain. While studies have shown that SEMG results have detected different pathologies in patients with back pain, none of the studies reported health outcomes. There are also no data on the impact of SEMG for managing back pain. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

### Additional Information

Not applicable.

## Related Policies

- N/A

## Benefit Application

Blue Shield of California Promise Health Plan is contracted with L.A. Care Health Plan for Los Angeles County and the Department of Health Care Services for San Diego County to provide Medi-Cal health benefits to its Medi-Cal recipients. In order to provide the best health care services and practices, Blue Shield of California Promise Health Plan has an extensive network of Medi-Cal primary care providers and specialists. Recognizing the rich diversity of its membership, our providers are given training and educational materials to assist in understanding the health needs of their patients as it could be affected by a member's cultural heritage.

The benefit designs associated with the Blue Shield of California Promise Medi-Cal plans are described in the Member Handbook (also called Evidence of Coverage).

## Regulatory Status

Surface electromyography devices approved by the U.S. Food and Drug Administration (FDA) include those that use a single electrode or a fixed array of multiple surface electrodes. Examples include the CMAP Pro (Medical Technologies) and Model 9200 EMG System (Myotronics-Noromed).

Several FDA approved devices combine SEMG along the spine with other types of monitors. For example, in 2007, the Insight Discovery (Fasstech) was cleared for marketing through the 510(k) process. The device contains 6 sensor types, 1 of which is for SEMG. The indications include measuring bilateral differences in SEMG along the spine and measuring SEMG along the spine during functional tasks. (Earlier Insight models had fewer sensors.) FDA product code: IKN.

## Health Equity Statement

Blue Shield of California Promise Health Plan's mission is to transform its health care delivery system into one that is worthy of families and friends. Blue Shield of California Promise Health Plan seeks to advance health equity in support of achieving Blue Shield of California Promise Health Plan's mission.

Blue Shield of California Promise Health Plan ensures all Covered Services are available and accessible to all members regardless of sex, race, color, religion, ancestry, national origin, ethnic group identification, age, mental disability, physical disability, medical condition, genetic information, marital status, gender, gender identity, or sexual orientation, or identification with any other persons or groups defined in Penal Code section 422.56, and that all Covered Services are provided in a culturally and linguistically appropriate manner.

## Rationale

### Background Back Pain

Back pain is a common condition that affects most individuals at some point in their lives.<sup>1</sup> Identifying the pathogenesis of back pain is challenging, in part due to the complex anatomy of the back, which includes vertebrae, intervertebral discs, facet joints, spinal nerve roots, and numerous muscles. Back pain may be related to osteoarthritis, disc disease, subluxation, or muscular pathologies, such as muscle strain or spasm. Moreover, due to referred pain patterns, the location of the pain may not be

anatomically related to the pathogenesis of the pain. For example, buttock or leg pain may be related to pathology in the spine. In addition to the diagnostic challenges of back pain is the natural history of acute back pain.

### Diagnosis

Aside from physical examination, diagnostic testing includes imaging technologies, such as magnetic resonance imaging, designed to identify pathology (e.g., bulging discs), or tests such as discography to localize the abnormality by reproducing the pain syndrome.<sup>1</sup> However, these tests lack specificity and must be carefully interpreted in the context of the clinical picture. For example, magnetic resonance imaging identifies 5% of asymptomatic patients as having bulging discs. However, the presence of a bulging disc may only be clinically significant if correlated with other symptoms. Assessment of the musculature may focus on a range of motion or strength exercises.

In contrast to anatomic imaging, surface electromyography (SEMG), which records the summation of muscle activity from groups of muscles, has been investigated as a technique to evaluate the physiologic functioning of the back.<sup>2</sup> A noninvasive procedure, SEMG differs from needle electromyography, an invasive procedure in which the electrical activity of individual muscles is recorded. Paraspinal SEMG has been explored to evaluate abnormal patterns of electrical activity in the paraspinal muscles in patients with back pain symptoms such as spasm, tenderness, limited range of motion, or postural disorders. The technique is performed using a single or an array of electrodes placed on the skin surface, with recordings made at rest, in various positions, or after a series of exercises. Recordings can also be made by using a handheld device, which is applied to the skin at different sites. Electrical activity is assessed by computer analysis of the frequency spectrum (i.e., spectral analysis), amplitude, or root mean square of the electrical action potentials. In particular, a spectral analysis that focuses on the median frequency has been used to assess paraspinal muscle fatigue during isometric endurance exercises. Paraspinal SEMG has been researched as a technique to establish the etiology of back pain and has been used to monitor the response to therapy and establish physical activity limits, such as assessing capacity to lift heavy objects or ability to return to work.

Paraspinal SEMG is an office-based procedure. The following clinical applications of the paraspinal SEMG have been proposed:

- clarification of diagnosis (i.e., muscle, joint, or disc disease)
- selection of a course of medical therapy
- selection of a type of physical therapy
- preoperative evaluation
- postoperative rehabilitation
- follow-up of acute low back pain (LBP)
- evaluation of exacerbation of chronic LBP
- evaluation of pain management treatment techniques.

### Treatment

Most cases of acute LBP resolve with conservative therapy (e.g., physical therapy) while continuing normal activities within limits permitted by the pain.<sup>1</sup> Therefore, initial imaging or other diagnostic testing is generally not recommended unless "red flag" warning signs are present or the pain persists for more than 4 to 6 weeks. Red flag findings include significant trauma, history of cancer, unrelenting night pain, fevers or chills, and progressive motor or sensory deficits.

### Literature Review

Evidence reviews assess whether a medical test is clinically useful. A useful test provides information to make a clinical management decision that improves the net health outcome. That is, the balance of benefits and harms is better when the test is used to manage the condition than when another test or no test is used to manage the condition.

The first step in assessing a medical test is to formulate the clinical context and purpose of the test. The test must be technically reliable, clinically valid, and clinically useful for that purpose. Evidence reviews assess the evidence on whether a test is clinically valid and clinically useful. Technical reliability is outside the scope of these reviews, and credible information on technical reliability is available from other sources.

### **Surface Electromyography**

Paraspinal surface electromyography (SEMG) has been used as a research tool to evaluate the performance of paraspinal muscles in patients with back pain and to further understand the etiology of low back pain (LBP).<sup>3,4,5,6</sup> Preliminary research has also been performed to determine which SEMG parameters best differentiate patients with and without back pain.<sup>7,8</sup>

### **Clinical Context and Test Purpose**

The purpose of paraspinal SEMG in individuals who have back pain is to identify the pathogenesis of the pain (i.e., muscle, joint, or disc disease) to inform a decision on a treatment plan.

The following PICO was used to select literature to inform this review.

#### ***Populations***

The relevant population of interest is individuals with back pain.

#### ***Interventions***

Paraspinal SEMG is a noninvasive technique that aggregates data on muscle activity from groups of muscles. One or more electrodes are placed on the skin surface, and recordings are taken at rest, in various positions, or during a series of exercises.

#### ***Comparators***

Other noninvasive techniques to assess back pain include clinical examination and imaging technologies.

#### ***Outcomes***

The general outcomes of interest are a reduction in back pain and improvement in activities of daily living.

Both false-positive test results and false-negative results can lead to an incorrect recommendation for the type of treatment or no treatment at all. Some treatments are long-term programs, and if individuals are incorrectly referred to the program, more appropriate therapy will be delayed.

### **Study Selection Criteria**

For the evaluation of clinical validity of the paraspinal SEMG test, studies that meet the following eligibility criteria were considered:

- Reported on the accuracy of the marketed version of the technology (including any algorithms used to calculate scores);
- Included a suitable reference standard;
- Patient/sample clinical characteristics were described;
- Patient/sample selection criteria were described.

### **Clinically Valid**

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

## Review of Evidence

No articles that directly compare the results of SEMG (which tests groups of muscles) with needle electromyography (which tests individual muscles) for diagnosing any specific muscle pathology were identified in literature searches. However, the pathology of individual muscles (i.e., radiculopathy, neuropathy) may represent a different process than the pathology of muscle groups (i.e., muscle strain, spasm), and thus SEMG may be considered by its advocates as a unique test for which there is currently no criterion standard. Nevertheless, even if one accepts this premise, there are inadequate data to evaluate the diagnostic performance of SEMG. In some instances, the asymmetrical electrical activity may have been used to define abnormality; results may be compared with normative data. However, no published literature was identified defining what degree of asymmetry would constitute abnormality.

A study by du Rose and Breen (2016) looked into the relationship between lumbar intervertebral range of motion and paraspinal muscle activity in healthy adults, as measured by SEMG and quantitative fluoroscopy, to establish "normal" measurements. Fluoroscopic images and SEMG measurements were taken for 20 men with no history of LBP. What would be considered normal intervertebral ranges of motion were related to a diverse set of muscle activation patterns as measured by SEMG. The authors concluded that larger sample sizes and measurements from patients with LBP would be needed to establish standard criterion.

Absent a criterion standard diagnostic test, correlation with the clinical symptoms and physical exam is critical. De Luca (1993) published a series of studies investigating a type of SEMG called the Back Analysis System, consisting of surface electrodes and other components to measure the electrical activity of muscles during isometric exercises designed to produce muscle fatigue.<sup>4</sup> Using physical exam and clinical history as a criterion standard, De Luca (1993) found that the Back Analysis System accurately identified control and back pain patients 84% and 91% of the time, respectively, with the values increasing to 100% in some populations. Accuracy was defined as the sum of true-positive and true-negative results. However, these studies were not designed as a clinical diagnostic tool per se but were intended to investigate the etiology of back pain and to investigate muscular fatigue patterns in patients with and without back pain.

Hu et al (2010, 2014) published 2 articles on dynamic topography, an approach to analyzing SEMG findings.<sup>10,11</sup> Both studies included patients with LBP and healthy controls. All participants underwent SEMG at study enrollment and then back pain patients participated in a rehabilitation program. The first study found different dynamic topography at baseline between the healthy people and back pain samples (a more symmetric pattern in healthy controls).<sup>10</sup> After physical therapy, the dynamic topography images of back pain patients were more similar to the healthy controls on some of the parameters assessed. In the second study, following rehabilitation, back pain patients were classified as responders or nonresponders based on changes in back pain severity.<sup>11</sup> Some associations were found between baseline SEMG parameters and response to rehabilitation. Surface electromyography was not repeated after the rehabilitation program, and thus it is unclear whether there are any significant associations between continued symptoms and SEMG abnormalities. Moreover, it is unclear how SEMG analysis would affect treatment decisions for patients with LBP.

## Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, more effective therapy, or avoid unnecessary therapy or testing.

## Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials.

A number of studies have described SEMG as an aid in classifying LBP.<sup>12,13,14,15,16</sup> Most of this research has focused on the use of SEMG to assess muscle fatigability rather than on how information from test findings could enhance patient management. While SEMG may be used to document muscle spasm or other muscular abnormalities objectively, it is unclear how such objective documentation would supplant or enhance clinical evaluation, or how this information would be used to alter the treatment plan. In part, the difficulty in clinical interpretation is understanding the extent to which the SEMG abnormalities are primary or secondary. Additionally, as noted in the Background section, no specific workup is recommended for acute LBP without warning signs.

The following studies have proposed using SEMG results to inform treatment decisions; however, none provided data to validate whether treatment based on SEMG results in improved outcomes.

In a study of patients with chronic LBP (N=216) by Kienbacher et al (2016), SEMG showed potential to discriminate between impaired and unimpaired neuromuscular regulation of back extensors, which would provide useful information for designing individualized exercise programs.<sup>17</sup>

In a study of patients with LBP (n=27) and pain-free controls (n=23) by Schabrun et al (2017), SEMG detected a loss of discrete motor cortical organization of the paraspinal muscles among those with LBP.<sup>18</sup> The invasive technique of needle electromyography is usually performed to detect this pathology. Patients with cortical reorganization may benefit from motor skill training.

In 2 older studies (1988, 1992), SEMG was shown to differentiate muscle spasm from muscle contracture.<sup>19,20</sup> Muscle spasm would be treated with relaxation therapy, and contracture would be treated with stretching exercises.

### **Chain of Evidence**

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility. Current evidence on clinical validity does not permit construction of a chain of evidence to support the use of SEMG as a diagnostic tool for evaluating and monitoring back pain.

### **Supplemental Information**

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

### **Practice Guidelines and Position Statements**

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a U.S. professional society, an international society with U.S. representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

### **American College of Occupational and Environmental Medicine**

In 2019, the guideline from the American College of Occupational and Environmental Medicine on diagnostic tests for low back disorders does not recommend surface electromyography as a technique for diagnosing low back disorders, based on insufficient evidence of efficacy.<sup>2</sup>

### **North American Spine Society and American Academy of Pain Medicine**

In 2020, the North American Spine Society with input from the American Academy of Pain Medicine issued a guideline on the diagnosis and treatment of low back pain.<sup>21</sup> When discussing the diagnostic accuracy of nonimaging tests, the guideline lacks any statement on surface electromyography.

### **U.S. Preventive Services Task Force Recommendations**

Not applicable.

### Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

### Ongoing and Unpublished Clinical Trials

A search of ClinicalTrials.gov in April 2025 did not identify any ongoing or unpublished trials that would likely influence this review.

## References

1. Knezevic NN, Candido KD, Vlaeyen JWS, et al. Low back pain. *Lancet*. Jul 03 2021; 398(10294): 78-92. PMID 34115979
2. Hegmann KT, Travis R, Belcourt RM, et al. Diagnostic Tests for Low Back Disorders. *J Occup Environ Med*. Apr 2019; 61(4): e155-e168. PMID 30694882
3. Cram JR, Lloyd J, Cahn TS. The reliability of EMG muscle scanning. *Int J Psychosom*. 1994; 41(1-4): 41-5. PMID 7843866
4. De Luca CJ. Use of the surface EMG signal for performance evaluation of back muscles. *Muscle Nerve*. Feb 1993; 16(2): 210-6. PMID 8429847
5. Jones SL, Hitt JR, DeSarno MJ, et al. Individuals with non-specific low back pain in an active episode demonstrate temporally altered torque responses and direction-specific enhanced muscle activity following unexpected balance perturbations. *Exp Brain Res*. Sep 2012; 221(4): 413-26. PMID 22875027
6. Sheeran L, Sparkes V, Caterson B, et al. Spinal position sense and trunk muscle activity during sitting and standing in nonspecific chronic low back pain: classification analysis. *Spine (Phila Pa 1976)*. Apr 15 2012; 37(8): E486-95. PMID 22024899
7. Hanada EY, Johnson M, Hubley-Kozey C. A comparison of trunk muscle activation amplitudes during gait in older adults with and without chronic low back pain. *PM R*. Oct 2011; 3(10): 920-8. PMID 22024323
8. Neblett R, Brede E, Mayer TG, et al. What is the best surface EMG measure of lumbar flexion-relaxation for distinguishing chronic low back pain patients from pain-free controls?. *Clin J Pain*. Apr 2013; 29(4): 334-40. PMID 23328325
9. du Rose A, Breen A. Relationships between Paraspinal Muscle Activity and Lumbar Inter-Vertebral Range of Motion. *Healthcare (Basel)*. Jan 05 2016; 4(1). PMID 27417592
10. Hu Y, Siu SH, Mak JN, et al. Lumbar muscle electromyographic dynamic topography during flexion-extension. *J Electromyogr Kinesiol*. Apr 2010; 20(2): 246-55. PMID 19540776
11. Hu Y, Kwok JW, Tse JY, et al. Time-varying surface electromyography topography as a prognostic tool for chronic low back pain rehabilitation. *Spine J*. Jun 01 2014; 14(6): 1049-56. PMID 24530438
12. Hung CC, Shen TW, Liang CC, et al. Using surface electromyography (SEMG) to classify low back pain based on lifting capacity evaluation with principal component analysis neural network method. *Annu Int Conf IEEE Eng Med Biol Soc*. 2014; 2014: 18-21. PMID 25569886
13. Humphrey AR, Nargol AV, Jones AP, et al. The value of electromyography of the lumbar paraspinal muscles in discriminating between chronic-low-back-pain sufferers and normal subjects. *Eur Spine J*. Mar 2005; 14(2): 175-84. PMID 15549487
14. Peach JP, McGill SM. Classification of low back pain with the use of spectral electromyogram parameters. *Spine (Phila Pa 1976)*. May 15 1998; 23(10): 1117-23. PMID 9615362
15. Roy SH, Oddsson LI. Classification of paraspinal muscle impairments by surface electromyography. *Phys Ther*. Aug 1998; 78(8): 838-51. PMID 9711209
16. Van Damme B, Stevens V, Perneel C, et al. A surface electromyography based objective method to identify patients with nonspecific chronic low back pain, presenting a flexion related movement control impairment. *J Electromyogr Kinesiol*. Dec 2014; 24(6): 954-64. PMID 25304196

17. Kienbacher T, Fehrmann E, Habenicht R, et al. Age and gender related neuromuscular pattern during trunk flexion-extension in chronic low back pain patients. J Neuroeng Rehabil. Feb 19 2016; 13: 16. PMID 26896325
18. Schabrun SM, Elgueta-Cancino EL, Hodges PW. Smudging of the Motor Cortex Is Related to the Severity of Low Back Pain. Spine (Phila Pa 1976). Aug 01 2017; 42(15): 1172-1178. PMID 25893342
19. Ellestad SM, Nagle RV, Boesler DR, et al. Electromyographic and skin resistance responses to osteopathic manipulative treatment for low-back pain. J Am Osteopath Assoc. Aug 1988; 88(8): 991-7. PMID 2975645
20. Bittman B, Cram JR. Surface electromyography: an electrophysiological alternative in pain management. Paper presented at: Presented at the American Pain Society; Oct 22-25 1992; San Diego, CA.
21. Kreiner DS, Matz P, Bono CM, et al. Guideline summary review: an evidence-based clinical guideline for the diagnosis and treatment of low back pain. Spine J. Jul 2020; 20(7): 998-1024. PMID 32333996

## Documentation for Clinical Review

### Please provide the following documentation:

- History and physical and/or consultation notes including:
  - Clinical findings (i.e., pertinent symptoms and duration)
  - Comorbidities
  - Activity and functional limitations
  - Reason for procedure/test/device, when applicable
  - Pertinent past procedural and surgical history
  - Past and present diagnostic testing and results
  - Prior conservative treatments, duration, and response
  - Treatment plan (i.e., surgical intervention)
- Consultation and medical clearance report(s), when applicable
- Radiology report(s) and interpretation (i.e., MRI, CT, discogram)
- Laboratory results
- Other pertinent multidisciplinary notes/reports: (i.e., psychological or psychiatric evaluation, physical therapy, multidisciplinary pain management), when applicable

### Post Service (in addition to the above, please include the following):

- Results/reports of tests performed
- Procedure report(s)

## Coding

*The list of codes in this Medical Policy is intended as a general reference and may not cover all codes. Inclusion or exclusion of a code(s) does not constitute or imply member coverage or provider reimbursement policy.*

Type	Code	Description
CPT®	95999	Unlisted neurological or neuromuscular diagnostic procedure
	96002	Dynamic surface electromyography, during walking or other functional activities, 1-12 muscles
	97799	Unlisted physical medicine/rehabilitation service or procedure
HCPCS	S3900	Surface electromyography (EMG)



## Policy History

This section provides a chronological history of the activities, updates and changes that have occurred with this Medical Policy.

Effective Date	Action
12/01/2025	New policy.

## Definitions of Decision Determinations

**Healthcare Services:** For the purpose of this Medical Policy, Healthcare Services means procedures, treatments, supplies, devices, and equipment.

**Medically Necessary or Medical Necessity** means reasonable and necessary services to protect life, to prevent significant illness or significant disability, or alleviate severe pain through the diagnosis or treatment of disease, illness, or injury, as required under W&I section 14059.5(a) and 22 CCR section 51303(a). Medically Necessary services must include services necessary to achieve age-appropriate growth and development, and attain, maintain, or regain functional capacity.

For Members less than 21 years of age, a service is Medically Necessary if it meets the Early and Periodic Screening, Diagnostic, and Treatment (EPSDT) standard of Medical Necessity set forth in 42 USC section 1396d(r)(5), as required by W&I sections 14059.5(b) and 14132(v). Without limitation, Medically Necessary services for Members less than 21 years of age include all services necessary to achieve or maintain age-appropriate growth and development, attain, regain or maintain functional capacity, or improve, support, or maintain the Member's current health condition. Contractor must determine Medical Necessity on a case-by-case basis, taking into account the individual needs of the Child.

### Criteria Determining Experimental/Investigational Status

In making a determination that any procedure, treatment, therapy, drug, biological product, facility, equipment, device, or supply is "experimental or investigational" by the Plan, the Plan shall refer to evidence from the national medical community, which may include one or more of the following sources:

1. Evidence from national medical organizations, such as the National Centers of Health Service Research.
2. Peer-reviewed medical and scientific literature.
3. Publications from organizations, such as the American Medical Association (AMA).
4. Professionals, specialists, and experts.
5. Written protocols and consent forms used by the proposed treating facility or other facility administering substantially the same drug, device, or medical treatment.
6. An expert physician panel selected by one of two organizations, the Managed Care Ombudsman Program of the Medical Care Management Corporation or the Department of Managed Health Care.

## Feedback

Blue Shield of California Promise Health Plan is interested in receiving feedback relative to developing, adopting, and reviewing criteria for medical policy. Any licensed practitioner who is contracted with Blue Shield of California Promise Health Plan is welcome to provide comments, suggestions, or concerns. Our internal policy committees will receive and take your comments into consideration. Our medical policies are available to view or download at [www.blueshieldca.com/en/bsp/providers](http://www.blueshieldca.com/en/bsp/providers).

For medical policy feedback, please send comments to: [MedPolicy@blueshieldca.com](mailto:MedPolicy@blueshieldca.com)

Questions regarding the applicability of this policy should be directed to the Blue Shield of California Promise Health Plan Prior Authorization Department at (800) 468-9935, or the Complex Case Management Department at (855) 699-5557 (TTY 711) for San Diego County and (800) 605-2556 (TTY 711) for Los Angeles County or visit the provider portal at [www.blueshieldca.com/en/bsp/providers](http://www.blueshieldca.com/en/bsp/providers).

*Disclaimer: Blue Shield of California Promise Health Plan may consider published peer-reviewed scientific literature, national guidelines, and local standards of practice in developing its medical policy. Federal and state law, as well as member health services contract language, including definitions and specific contract provisions/exclusions, take precedence over medical policy and must be considered first in determining covered services. Member health services contracts may differ in their benefits. Blue Shield of California Promise Health Plan reserves the right to review and update policies as appropriate.*